arXiv:1411.2628v1 [q-fin.CP] 10 Nov 2014

Exact solution of a generalized version of the Black-Scholes equation

L.-A. Cotfas[a,*], C. Delcea[a], N. Cotfas[b]

[a] *Faculty of Economic Cybernetics, Statistics and Informatics, Bucharest University of Economic Studies, 6 Piata Romana, 010374 Bucharest, Romania*
[b] *University of Bucharest, Physics Department, P.O. Box MG-11, 077125 Bucharest, Romania*

Abstract

We analyze a generalized version of the Black-Scholes equation depending on a parameter $a \in (-\infty, 0)$. It satisfies the martingale condition and coincides with the Black-Scholes equation in the limit case $a \nearrow 0$. We show that the generalized equation is exactly solvable in terms of Hermite polynomials and numerically compare its solution with the solution of the Black-Scholes equation.

Keywords: econophysics, quantum finance, Black-Scholes equation, option pricing
2010 MSC: 91B80, 91G80

1. Introduction

The mathematical model based on the Black-Scholes equation

$$\frac{\partial C}{\partial t} = -\frac{\sigma^2}{2} S^2 \frac{\partial^2 C}{\partial S^2} - rS\frac{\partial C}{\partial S} + rC \tag{1}$$

anticipates rather well the observed prices for options in the case of a strike price that is not too far from the current price of the underlying asset [8]. The price of an option at a moment of time t depends on the current price S, the volatility σ, the risk-free interest rate r, the strike price K and the

*Corresponding author
Email addresses: lcotfas@gmail.com (L.-A. Cotfas), camelia.delcea@yahoo.com (C. Delcea), ncotfas@yahoo.com (N. Cotfas)

maturity time T. In the case of an European option, the price is described by the solution $C(S,t)$ of equation (1) satisfying the condition

$$C(S,T) = \begin{cases} S-K & \text{if} \quad S \geq K \\ 0 & \text{if} \quad S < K \end{cases} \tag{2}$$

in the case of a call option, and

$$C(S,T) = \begin{cases} 0 & \text{if} \quad S \geq K \\ K-S & \text{if} \quad S < K \end{cases} \tag{3}$$

in the case of a put option. The alternative version of the equation (1)

$$\frac{\partial \tilde{C}}{\partial t} = -\frac{\sigma^2}{2}\frac{\partial^2 \tilde{C}}{\partial x^2} + \left(\frac{\sigma^2}{2} - r\right)\frac{\partial \tilde{C}}{\partial x} + r\tilde{C} \tag{4}$$

obtained by using the change of independent variable

$$S = \mathrm{e}^x \tag{5}$$

allows one to use the formalism of quantum mechanics in option pricing [1, 2, 4, 7]. In the new variable the conditions (2) and (3) become

$$\tilde{C}(x,T) = \begin{cases} \mathrm{e}^x - K & \text{if} \quad x \geq \ln K \\ 0 & \text{if} \quad x < \ln K \end{cases} \tag{6}$$

and respectively,

$$\tilde{C}(x,T) = \begin{cases} 0 & \text{if} \quad x \geq \ln K \\ K - \mathrm{e}^x & \text{if} \quad x < \ln K\,. \end{cases} \tag{7}$$

The more general version of (4) depending on a function $V(x)$

$$\frac{\partial \tilde{C}}{\partial t} = -\frac{\sigma^2}{2}\frac{\partial^2 \tilde{C}}{\partial x^2} + \left(\frac{\sigma^2}{2} - V(x)\right)\frac{\partial \tilde{C}}{\partial x} + V(x)\,\tilde{C} \tag{8}$$

satisfies the martingale condition [1] and hence can be used for studying processes in finance. Our purpose is to investigate the particular case

$$V(x) = ax + r \tag{9}$$

that is, the equation

$$\frac{\partial \tilde{C}}{\partial t} = -\frac{\sigma^2}{2}\frac{\partial^2 \tilde{C}}{\partial x^2} + \left(\frac{\sigma^2}{2} - ax - r\right)\frac{\partial \tilde{C}}{\partial x} + (ax+r)\tilde{C} \tag{10}$$

where $a \in (-\infty, 0)$ is a parameter. The equation (10) is exactly solvable in terms of Hermite polynomials and coincides in the limit case $a \nearrow 0$ with the equation (4) which corresponds to the standard Black-Scholes equation.

2. A shifted oscillator

Let $\alpha \in (-\infty, 0)$ and $\beta \in \mathbb{R}$ be two constants. By using the Hermite polynomial

$$\mathbf{H}_n(s) = (-1)^n \, \mathrm{e}^{s^2} \, \frac{d^n}{ds^n}(\mathrm{e}^{-s^2}) \tag{11}$$

we define for each $n \in \{0, 1, 2, ...\}$ the function

$$\psi_n(x) = \tfrac{1}{\sqrt{n!\, 2^n}} \sqrt[4]{\tfrac{-\alpha}{2\pi}} \, \mathrm{e}^{\frac{\alpha}{4}x^2 + \frac{\beta}{2}x + \frac{\beta^2}{4\alpha}} \, \mathbf{H}_n\left(\sqrt{\tfrac{-\alpha}{2}}x - \tfrac{\beta}{\sqrt{-2\alpha}}\right). \tag{12}$$

If we denote

$$s = \sqrt{\tfrac{-\alpha}{2}}x - \tfrac{\beta}{\sqrt{-2\alpha}} \tag{13}$$

then the previous relation can be written as

$$\mathbf{H}_n(s) = \sqrt{n!\, 2^n} \sqrt[4]{\tfrac{2\pi}{-\alpha}} \, \mathrm{e}^{\frac{1}{2}s^2} \, \psi_n\left(\sqrt{\tfrac{2}{-\alpha}}s - \tfrac{\beta}{\alpha}\right). \tag{14}$$

By substituting this relation into the diferential equation

$$\mathbf{H}_n''(s) - 2s\mathbf{H}_n'(s) + 2n\mathbf{H}_n(s) = 0 \tag{15}$$

satisfied by the Hermite polynomial $\mathbf{H}_n$ we get the equality

$$-\psi_n''\left(\sqrt{\tfrac{2}{-\alpha}}s - \tfrac{\beta}{\alpha}\right) + \left(-\tfrac{\alpha}{2}s^2 + \tfrac{\alpha}{2} + \alpha n\right) \, \psi_n\left(\sqrt{\tfrac{2}{-\alpha}}s - \tfrac{\beta}{\alpha}\right) = 0 \tag{16}$$

which can be written in the form

$$\left(-\frac{\partial^2}{\partial x^2} + \frac{(\alpha x+\beta)^2}{4} + \frac{\alpha}{2}\right)\psi_n = -\alpha n \, \psi_n. \tag{17}$$

This means that ψ_n is an eigenfunction of the shifted oscillator [3, 5, 6].

$$H = -\frac{\partial^2}{\partial x^2} + \frac{(\alpha x+\beta)^2}{4} + \frac{\alpha}{2} \tag{18}$$

corresponding to the eigenvalue $\lambda_n = -\alpha n$, for any $n \in \{0, 1, 2, ...\}$. Since

$$\int_{-\infty}^{\infty} \mathrm{e}^{-s^2}\ \mathbf{H}_n(s)\ \mathbf{H}_k(s)\ ds = \begin{cases} n!\, 2^n \sqrt{\pi} & \text{if} \quad n = k \\ 0 & \text{if} \quad n \neq k \end{cases} \tag{19}$$

the system of functions $\{\psi_n\}_{n=0,1,2,...}$ is orthonormal, that is,

$$\int_{-\infty}^{\infty} \psi_n(x)\ \psi_k(x)\ dx = \begin{cases} 1 & \text{if} \quad n = k \\ 0 & \text{if} \quad n \neq k. \end{cases} \tag{20}$$

One can prove that it is complete in the space of square integrable functions.

3. A generalized version of the Black-Scholes equation

A straightforward generalization of the equation

$$\frac{\partial \tilde{C}}{\partial t} = -\frac{\sigma^2}{2} S^2 \frac{\partial^2 \tilde{C}}{\partial S^2} - rS\frac{\partial \tilde{C}}{\partial S} + r\tilde{C} \tag{21}$$

is the equation

$$\frac{\partial \tilde{C}}{\partial t} = -\frac{\sigma^2}{2}\frac{\partial^2 \tilde{C}}{\partial x^2} + \left(\frac{\sigma^2}{2} - V(x)\right)\frac{\partial \tilde{C}}{\partial x} + V(x)\,\tilde{C} \tag{22}$$

satisfying the martingale condition [1]. It can be written in the form

$$\frac{\partial \tilde{C}}{\partial t} = H_V \tilde{C} \tag{23}$$

by using the Hamiltonian

$$H_V = -\frac{\sigma^2}{2}\frac{\partial^2}{\partial x^2} + \left(\frac{\sigma^2}{2} - V(x)\right)\frac{\partial}{\partial x} + V(x). \tag{24}$$

The Hamiltonian H_V is equivalent with the Hermitian Hamiltonian

$$H_{\text{eff}} = -\frac{\sigma^2}{2}\frac{\partial^2}{\partial x^2} + \frac{1}{2}\frac{\partial V}{\partial x} + \frac{1}{2\sigma^2}V^2 + \frac{1}{2}V + \frac{\sigma^2}{8} \tag{25}$$

by the similarity transformation

$$H_{\text{eff}} = \mathrm{e}^{-u} H_V \mathrm{e}^{u} \tag{26}$$

where

$$u = \frac{1}{2}x - \frac{1}{\sigma^2}\int_0^x V(y)\,dy. \tag{27}$$

In the particular case $V(x)=ax+r$ considered in this article

$$\begin{aligned} H_V &= -\tfrac{\sigma^2}{2}\tfrac{\partial^2}{\partial x^2} + \left(\tfrac{\sigma^2}{2} - ax - r\right)\tfrac{\partial}{\partial x} + ax + r \\ H_{\text{eff}} &= \tfrac{\sigma^2}{2}\left[-\tfrac{\partial^2}{\partial x^2} + \tfrac{1}{4}\left(\tfrac{2a}{\sigma^2}x + \tfrac{2r}{\sigma^2} + 1\right)^2 + \tfrac{a}{\sigma^2}\right] \\ u &= -\tfrac{a}{2\sigma^2}x^2 - \left(\tfrac{r}{\sigma^2} - \tfrac{1}{2}\right)x. \end{aligned} \tag{28}$$

The Hamiltonian H_{eff} is up to the multiplicative factor $\frac{\sigma^2}{2}$ the Hamiltonian of a shifted oscillator, namely

$$H_{\text{eff}} = \frac{\sigma^2}{2} H \tag{29}$$

where

$$H = -\tfrac{\partial^2}{\partial x^2} + \tfrac{(\alpha x+\beta)^2}{4} + \tfrac{\alpha}{2} \tag{30}$$

with

$$\alpha = \tfrac{2a}{\sigma^2} \qquad \text{and} \qquad \beta = \tfrac{2r}{\sigma^2}+1.$$

Therefore, for each $n \in \{0, 1, 2, ...\}$ the function

$$\psi_n(x) = \tfrac{1}{\sqrt{n!\,2^n\sigma}}\sqrt[4]{\tfrac{-a}{\pi}}\;\mathrm{e}^{\frac{a}{2\sigma^2}x^2 + \left(\frac{r}{\sigma^2}+\frac{1}{2}\right)x + \frac{\sigma^2}{8a}\left(\frac{2r}{\sigma^2}+1\right)^2}\mathbf{H}_n\left(\tfrac{\sqrt{-a}}{\sigma}x - \tfrac{\sigma}{2\sqrt{-a}}\left(\tfrac{2r}{\sigma^2}+1\right)\right) \tag{31}$$

is an eigenfunction of H_{eff} corresponding to the eigenvalue $-an$

$$H_{\text{eff}}\,\psi_n = -an\,\psi_n. \tag{32}$$

The equation (23) can be written as

$$\frac{\partial \tilde{C}}{\partial t} = \mathrm{e}^{u} H_{\text{eff}} \mathrm{e}^{-u}\,\tilde{C} \tag{33}$$

or in the form

$$\frac{\partial}{\partial t}\,\mathrm{e}^{-u}\tilde{C} = H_{\text{eff}}\,\mathrm{e}^{-u}\tilde{C}. \tag{34}$$

A function $\tilde{C}(x,t)$ is the solution of (23) satisfying (6) if and only if

$$\psi(x,t) = \mathrm{e}^{-u}\tilde{C}(x,t) \tag{35}$$

that is, the function

$$\psi(x,t) = \mathrm{e}^{\frac{a}{2\sigma^2}x^2+\left(\frac{r}{\sigma^2}-\frac{1}{2}\right)x}\tilde{C}(x,t) \tag{36}$$

is the solution of the equation

$$\frac{\partial\psi}{\partial t} = H_{\mathrm{eff}}\,\psi \tag{37}$$

satisfying the condition

$$\psi(x,T) = \mathrm{e}^{\frac{a}{2\sigma^2}x^2+\left(\frac{r}{\sigma^2}-\frac{1}{2}\right)x}\begin{cases} \mathrm{e}^x-K & \text{if} \quad x\geq\ln K \\ 0 & \text{if} \quad x<\ln K\,. \end{cases} \tag{38}$$

But, the solution of (37) satisfying (38) is

$$\psi(x,t) = \sum_{n=0}^{\infty} c_n\,\mathrm{e}^{-ant}\;\psi_n(x) \tag{39}$$

with the coefficients c_n determined from the relation

$$\sum_{n=0}^{\infty} c_n\,\mathrm{e}^{-anT}\;\psi_n(x) = \mathrm{e}^{\frac{a}{2\sigma^2}x^2+\left(\frac{r}{\sigma^2}-\frac{1}{2}\right)x}\begin{cases} \mathrm{e}^x-K & \text{if} \quad x\geq\ln K \\ 0 & \text{if} \quad x<\ln K \end{cases} \tag{40}$$

namely,

$$c_n = \mathrm{e}^{anT}\int_{\ln K}^{\infty}\mathrm{e}^{\frac{a}{2\sigma^2}x^2+\left(\frac{r}{\sigma^2}-\frac{1}{2}\right)x}\,(\mathrm{e}^x-K)\psi_n(x)\,dx. \tag{41}$$

In the case of the call option, the solution of the generalized Black-Scholes equation expressed in terms of Hermite polynomials is

$$C_a(S,t) = \frac{S}{\sqrt{\sigma}}\sqrt[4]{\frac{-a}{\pi}}\mathrm{e}^{\frac{\sigma^2}{8a}\left(\frac{2r}{\sigma^2}+1\right)^2}\sum_{n=0}^{\infty} c_n\frac{\mathrm{e}^{-ant}}{\sqrt{n!\,2^n}}\,\mathbf{H}_n\!\left(\frac{\sqrt{-a}}{\sigma}\ln S-\frac{\sigma}{2\sqrt{-a}}\left(\frac{2r}{\sigma^2}+1\right)\right). \tag{42}$$

The case of a put option can be analyzed in a very similar way.

The Black-Scholes equation is exactly solvable. By denoting

$$\Phi(\zeta) = \frac{1}{\sqrt{2\pi}}\int_{-\infty}^{\zeta}\mathrm{e}^{-\eta/2}d\eta = \frac{1}{2}\left(1+\mathrm{erf}(x/\sqrt{2})\right) \tag{43}$$

the formulas for the values of a European option can be written in the form

$$\begin{aligned} C_{\rm call}(S,t) &= S\,\Phi(d_1) - K\,{\rm e}^{-r(T-t)}\,\Phi(d_2) \\ C_{\rm put}(S,t) &= K\,{\rm e}^{-r(T-t)}\,\Phi(d_1) - S\,\Phi(d_2) \end{aligned} \tag{44}$$

where [8]

$$d_1 = \frac{\ln(S/K)+\left(r+\frac{1}{2}\sigma^2\right)(T-t)}{\sigma\sqrt{T-t}}, \qquad d_2 = \frac{\ln(S/K)+\left(r-\frac{1}{2}\sigma^2\right)(T-t)}{\sigma\sqrt{T-t}}. \tag{45}$$

In Fig. 1 we present $C_a(S,t)$ (thick line) versus $C_{\rm call}(S,t)$ (thin line) and $C_{\rm call}(S,T)$ (dashed line) for $t=3$ (left side), $t=4$ (right side) and $a=-0.03$ (first row), $a=-0.02$ (second row), $a=-0.01$ (third row) by choosing the following values of the parameters: $\sigma=0.25$, $r=0.03$, $K=3$ and $T=5$.

4. Concluding remarks

In quantum mechanics as well as in econophysics, exact solutions are known only in a small number of particular cases and, generally, they play an important role. The generalized version of the Black-Scholes equation investigated in this article:

- is exactly solvable for any $a\in(-\infty,0)$,
- satisfies the martingale condition for any $a\in(-\infty,0)$,
- coincides with the Black-Scholes equation in the limit case $a \nearrow 0$.

In practice, there are some deviations of prices from those described by the solution of the Black-Scholes equation. We think that the solution of the generalized Black-Scholes equation might describe some observed prices, and the parameter a might have a certain financial meaning.

References

[1] B.E. Baaquie, Quantum Finance, Cambridge University Press, 2004.

[2] F. Bagarello, A quantum statistical approach to simplified stock markets, Physics A 388 (2009) 4397.

[3] F. Cooper, A. Khare, U. Sukhatme, Supersymmetry and quantum mechanics, Phys. Rep. 251 (1995) 267.

[4] L.-A. Cotfas, A finite-dimensional quantum model for the stock market, Physics A 392 (2013) 371.

[5] N. Cotfas and L.A. Cotfas, Hypergeometric type operators and their supersymmetric partners, J. Math. Phys. 52 (2011) 052101.

[6] M.A. Jafarizadeh and H. Fakhri, Parasupersymmetry and shape invariance in differential equations of mathematical physics and quantum mechanics, Ann. Phys. NY 262 (1998) 260.

[7] T.K. Jana and P. Roy, Supersymmetry in option pricing, Physica A 390 (2011) 2350-55.

[8] Ö. Uğur, An Introduction to Computational Finance, Imperial College Press, London, 2009.



Figure 1: The solution $C_a(S,t)$ (thick line) of the generalized Black-Scholes equation versus the solution $C_{\text{call}}(S,t)$ (thin line) of the Black-Scholes equation and the payoff function $C_{\text{call}}(S,T)$ (dashed line) for $t = 3$ (left hand side), $t = 4$ (right hand side) and $a = -0.03$ (first row), $a = -0.02$ (second row), $a = -0.01$ (third row).